

Vectors

A vector is a set of coordinates. Notation: \mathbf{v} or \vec{v} .

$$\vec{v} = \langle 1, 2, 3 \rangle$$

$$\vec{w} = \langle 1, 2 \rangle$$

Here, \vec{v} is a vector in \mathbb{R}^3 , and \vec{w} is a vector in \mathbb{R}^2 . The magnitude, or norm of a vector, represented by $\|\vec{v}\|$, is defined as $\sqrt{v_x^2 + v_y^2}$ in 2-space or $\sqrt{v_x^2 + v_y^2 + v_z^2}$ in 3-space.

Vector addition and subtraction

We can add vectors component-wise:

$$\vec{v} = \langle 1, 2, 3 \rangle$$

$$\vec{w} = \langle 4, 5, 6 \rangle$$

$$\vec{v} + \vec{w} = \langle 5, 7, 9 \rangle$$

We can also subtract vectors component-wise:

$$\vec{v} = \langle 4, 5, 6 \rangle$$

$$\vec{w} = \langle 1, 2, 3 \rangle$$

$$\vec{v} - \vec{w} = \langle 4 - 1, 5 - 2, 6 - 3 \rangle = \langle 3, 3, 3 \rangle$$

Geometrically, vector addition works by putting vectors "tip to tail."

Unit vectors

Vectors are often defined in terms of *unit vectors*:

In \mathbb{R}^2 :

$$\hat{i} = \langle 1, 0 \rangle$$

$$\hat{j} = \langle 0, 1 \rangle$$

In \mathbb{R}^3 :

$$\hat{i} = \langle 1, 0, 0 \rangle$$

$$\hat{j} = \langle 0, 1, 0 \rangle$$

$$\hat{k} = \langle 0, 0, 1 \rangle$$

For example:

$$\langle 1, 2, 3 \rangle = \hat{i} + 2\hat{j} + 3\hat{k}$$

Scalar multiplication

Vectors can be multiplied by scalars component-wise:

$$\lambda \langle a, b, c \rangle = \langle \lambda a, \lambda b, \lambda c \rangle$$

Dot products

Taking the dot product is a method of multiplying vectors to produce a scalar. The formula for a dot product is

$$\langle a, b \rangle \cdot \langle x, y \rangle = ax + by$$

$$\langle a, b, c \rangle \cdot \langle x, y, z \rangle = ax + by + cz$$

Another way to write this is:

$$\vec{a} \cdot \vec{b} = \|\vec{a}\| \|\vec{b}\| \cos(\theta)$$

Where θ is the angle between the vectors.

The dot product geometrically represents the scalar projection of one vector onto another.

Cross products

Taking the cross product is a method of multiplying vectors to produce a vector. The formula for a cross product is:

$$\langle a, b, c \rangle \times \langle x, y, z \rangle = \langle -cy + bz, cx - az, -bx + ay \rangle$$

Cross products are non-commutative. Order does matter. $\vec{a} \times \vec{b} \neq \vec{b} \times \vec{a}$ (except in some very specific circumstances).

Cross products geometrically:

- From a right hand system (i.e. \vec{a} , \vec{b} , and $\vec{a} \times \vec{b}$ form a right hand system)
- Are orthogonal to the crossed vectors
- Have magnitude equal to the parallelogram spanned by the crossed vectors

Cross products always have the magnitude:

$$\|\vec{a} \times \vec{b}\| = \|\vec{a}\| \|\vec{b}\| \sin(\theta)$$

Parallelepipeds

A *parallelepiped* is the shape spanned by three non-zero vectors. Given vectors \vec{u} , \vec{v} , and \vec{w} , the base of the parallelepiped is the parallelogram spanned by \vec{v} and \vec{w} . Therefore, the area of the base is $\|\vec{v} \times \vec{w}\|$. The volume of a parallelepiped \mathcal{D} spanned by \vec{u} , \vec{v} and \vec{w} can be calculated with the *scalar triple product*:

$$V(\mathcal{D}) = |\vec{u} \cdot (\vec{v} \times \vec{w})| = \det(u, v, w)$$

Lines

Lines are defined in terms of a point and a direction, or two points. Given a direction vector (parallel to the line), \vec{d} , and a point on the line \vec{p} , the parameterization of the line is:

$$\vec{r}(t) = \vec{p} + t\vec{d}$$

Given two points, \vec{p}_1 and \vec{p}_2 :

$$\vec{r}(t) = \vec{p}_1 + t(\vec{p}_2 - \vec{p}_1)$$

Planes

Planes are defined in terms of three non-colinear points, or a normal vector and a plane.

To get a normal vector from 3 points, A , B , and C , compute $\vec{AB} \times \vec{AC}$

With normal vector \vec{n} and point P :

$$\vec{n}_x(x - P_x) + \vec{n}_y(y - P_y) + \vec{n}_z(z - P_z) = 0$$

Polar Coordinates

Polar coordinates are used to represent points in \mathbb{R}^2 . They are represented as (r, θ) , where $r \in \mathbb{R}$ and $\theta \in [0, 2\pi)$.

To convert between Cartesian coordinates and polar coordinates:

$$\begin{aligned} r &= \sqrt{x^2 + y^2} & x &= r \cos \theta \\ \theta &= \arctan\left(\frac{y}{x}\right) & y &= r \sin \theta \end{aligned}$$

Mathematica snippet: `AngleVector[{x, y}]` will convert polar to rectangular.

Spherical Coordinates

Spherical coordinates are one of two generalizations to \mathbb{R}^3 of polar coordinates. They are represented as (ρ, θ, ϕ) , where $\rho \in \mathbb{R}$, $\theta \in [0, 2\pi)$, and $\phi \in [0, \pi]$. ρ represents the distance to the origin, θ represents the counterclockwise angle towards positive y in the xz -plane, and ϕ represents the angle towards positive x in the xy -plane.

$$\begin{aligned} \rho &= \sqrt{x^2 + y^2 + z^2} & x &= \rho \sin \phi \cos \theta \\ \theta &= \arctan\left(\frac{y}{x}\right) & y &= \rho \sin \phi \sin \theta \\ \rho &= \arccos\left(\frac{z}{\rho}\right) & z &= \rho \cos \phi \end{aligned}$$

Cylindrical Coordinates

Cylindrical coordinates are one of two generalizations to \mathbb{R}^3 of polar coordinates. They are represented as (r, θ, z) , where $r \in \mathbb{R}$, $\theta \in [0, 2\pi)$, and $z \in \mathbb{R}$. r represents the distance to the origin, θ represents the counterclockwise angle towards positive y in the xz -plane, and z represents the distance from the xy -plane in the positive z direction.

$$\begin{aligned} r &= \sqrt{x^2 + y^2 + z^2} & x &= r \cos \theta \\ \theta &= \arctan\left(\frac{y}{x}\right) & y &= r \sin \theta \\ z &= z & z &= z \end{aligned}$$

Surfaces to remember

Cylindrical Coordinates:	
equation	description
$r = R$	cylinder of radius R
$\theta = \theta_0$	vertical half-plane
$z = c$	horizontal plane
Spherical Coordinates:	
equation	description
$\rho = R$	sphere of radius R
$\theta = \theta_0$	vertical half-plane
$\phi = \phi_0$	right circular cone
Rectangular Coordinates:	
equation	description
$x^2 + y^2 = z^2$	right circular cone
$x^2 + y^2 + z^2 = R$	sphere (radius R)
$x^2 + y^2 = R$	cylinder (radius R)

Calculus of Vector-Valued Functions

Calculus can be done on vector-valued functions component-wise. This includes limits, differentiation, and integration. There are some additional differentiation rules

- Sum rule: $(\vec{r}_1(t) + \vec{r}_2(t))' = \vec{r}_1'(t) + \vec{r}_2'(t)$
 - Chain rule: $\vec{r}(g(t)) = g'(t)\vec{r}'(g(t))$
 - Product rules
 - Scalar product rule: $(\lambda \vec{r}(t))' = \lambda \vec{r}'(t)$
 - Dot product rule: $(\vec{r}_1 \cdot \vec{r}_2)' = \vec{r}_1' \cdot \vec{r}_2 + \vec{r}_1 \cdot \vec{r}_2'$
 - Cross product rule: $(\vec{r}_1 \times \vec{r}_2)' = \vec{r}_1' \times \vec{r}_2 + \vec{r}_1 \times \vec{r}_2'$
- Remember! Cross products are non-commutative.

The derivative of a vector is also called the *tangent vector*, or *velocity vector*. This is because if $\vec{r}'(t_0)$ is non-zero, it points in the direction tangent to the curve at $r(t_0)$. The tangent line has parametrization:

$$\vec{L}(t) = \vec{r}(t_0) + t\vec{r}'(t_0)$$

Arc length

If $\vec{r}(t) = \langle x(t), y(t), z(t) \rangle$ directly traverses curve \mathcal{L} , for $a \leq t \leq b$, the arc length, s of \mathcal{L} is:

$$\int_a^b \|\vec{r}'(t)\| dt = \int_a^b \sqrt{x'(t)^2 + y'(t)^2 + z'(t)^2} dt$$

Speed

The velocity vector, \vec{v} , points in the direction of travel. It's magnitude is the speed:

$$v(t) = \frac{ds}{dt} = \|\vec{r}'(t)\|$$

Functions in Multiple Variables

A function can exist in *multiple variables* if it takes several parameters.

If $f(x, y)$ is defined near $P = (a, b)$, then

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = L$$

if, $\forall \epsilon > 0 \exists \delta > 0$ s.t. $0 < d((x, y), (a, b)) < \delta$ then $|f(x, y) - L| < \epsilon$.

A function in two variables is *continuous* at (a, b) if the following equation holds:

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = f(a, b)$$

Partial Derivatives

The partial derivatives of $f(x, y)$ are defined as the limits:

$$f_x(a, b) = \left[\frac{\delta f}{\delta x} \right]_{(a,b)} = \lim_{h \rightarrow 0} \frac{f(a+h, b) - f(a, b)}{h}$$

$$f_y(a, b) = \left[\frac{\delta f}{\delta y} \right]_{(a,b)} = \lim_{k \rightarrow 0} \frac{f(a, b+k) - f(a, b)}{k}$$

Partial derivatives can be approximated for small Δx and Δy values:

$$f_x(a, b) \approx \frac{\Delta f}{\Delta x} = \frac{f(a + \Delta x, b) - f(a, b)}{\Delta x}$$

$$f_y(a, b) \approx \frac{\Delta f}{\Delta y} = \frac{f(a, b + \Delta y) - f(a, b)}{\Delta y}$$

The second-order partial derivatives are written as:

$$\frac{\delta^2}{\delta x^2} f = f_{xx} \quad \frac{\delta^2}{\delta x \delta y} f = f_{xy} \quad \frac{\delta^2}{\delta y^2} f = f_{yy}$$

Clairaut's Theorem

Clairaut's theorem states that mixed partials are equivalent as long as they are continuous: $f_{xy} = f_{yx}$.

Differentiability of Functions in Multiple Variables

$f(x, y)$ is *differentiable* at (a, b) if $f_x(a, b)$ and $f_y(a, b)$ both exist, and

$$\lim_{(x,y) \rightarrow (a,b)} \frac{f(x, y) - L(x, y)}{\sqrt{(x-a)^2 + (y-b)^2}} = 0$$

where

$$L(x, y) = f(a, b) + f_x(a, b) \times (x - a) + f_y(a, b) \times (y - b)$$

Gradient Vectors

The gradient vector is defined is

$$\mathbb{R}^2 : \nabla f = \left\langle \frac{\delta f}{\delta x}, \frac{\delta f}{\delta y} \right\rangle$$

$$\mathbb{R}^3 : \nabla f = \left\langle \frac{\delta f}{\delta x}, \frac{\delta f}{\delta y}, \frac{\delta f}{\delta z} \right\rangle$$

This gradient has the properties:

- ∇f_P points in the fastest rate of increase, and the rate of that increase is $||\nabla f_P||$
- $-\nabla f_P$ points in the fastest rate of decrease, and the rate of that decrease is $-||\nabla f_P||$
- $-\nabla f_P$ is orthogonal to level curves through P

The chain rule for paths also applies:

$$\frac{d}{dt} f(\vec{r}(t)) = \nabla f_{\vec{r}(t)} \cdot \vec{r}'(t)$$

Directional Derivatives

For unit vector $\vec{u} = \langle h, k \rangle$, $D_u f(a, b)$ is the directional derivative with respect to \vec{u} .

$$D_u f(a, b) = \lim_{t \rightarrow 0} \frac{f(a + th, b + tk) - f(a, b)}{t}$$

For a differentiable function f , the directional derivative in the direction \vec{u} is computed using the gradient:

$$D_u f(a, b) = \nabla f_{(a,b)} \cdot \vec{u} = ||\nabla f_{(a,b)}|| \cos \theta$$

Tangent Plane

The equation for a tangent plane can be thought of as the equation for a tangent line generalized into three dimensions. The equation for a tangent line at x_0 is (derived from point-slope form):

$$y = f(x_0) + f'(x_0) \times (x - x_0)$$

The equation for a tangent plane of $f(x, y)$ at the point (a, b) is:

$$z = f(a, b) + f_x(a, b) \times (x - a) + f_y(a, b) \times (y - b)$$

A normal vector can be computed using this tangent plane equation.

Linear Approximation

If $f(x, y)$ is differentiable at (a, b) , the *linearization* of f centered at (a, b) is:

$$L(x, y) = f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b)$$

You can use this linearization to compute *linear approximations* of functions:

$$f(a + \Delta x, b + \Delta y) \approx f(a, b) + f_x(a, b)\Delta x + f_y(a, b)\Delta y$$

$$\Delta f \approx f_x(a, b)\Delta x + f_y(a, b)\Delta y$$

This can be expressed in a differential form as well:

$$df = f_x(x, y)dx + f_y(x, y)dy = \frac{\delta f}{\delta x}dx + \frac{\delta f}{\delta y}dy$$

Chain rules

If $f(x, y, z)$ is a composite function (meaning that it can be represented in a form similar to $f(x(s, t), y(s, t), z(s, t))$, where s , and t are the *independent variables*), then

$$\frac{\delta f}{\delta s} = \frac{\delta f}{\delta x} \frac{\delta x}{\delta s} + \frac{\delta f}{\delta y} \frac{\delta y}{\delta s} + \frac{\delta f}{\delta z} \frac{\delta z}{\delta s}$$

$$\frac{\delta f}{\delta t} = \frac{\delta f}{\delta x} \frac{\delta x}{\delta t} + \frac{\delta f}{\delta y} \frac{\delta y}{\delta t} + \frac{\delta f}{\delta z} \frac{\delta z}{\delta t}$$

Critical Points

$P = (a, b)$ is a *critical point* if $f_x(a, b) = 0$ or $f_x(a, b)$ does not exist, and $f_y(a, b) = 0$ or $f_y(a, b)$ does not exist.

The *discriminant* of $f(x, y)$ at $P = (a, b)$ is defined as the quantity:

$$D(a, b) = f_{xx}(a, b) \times f_{yy}(a, b) - f_{xy}^2(a, b)$$

The *second derivative test* allows us to categorize critical points. Give a critical point, (a, b) on f :

$$D(a, b) > 0, f_{xx}(a, b) > 0 \implies \text{local minimum}$$

$$D(a, b) > 0, f_{xx}(a, b) < 0 \implies \text{local maximum}$$

$$D(a, b) < 0 \implies \text{saddle point}$$

$$D(a, b) = 0 \implies \text{test inconclusive}$$

To find extreme values, first find the values at the critical points, then compare to the maximum value along the boundary.

Lagrange Multipliers

If the local extreme of $f(x, y)$ is subject to a constraint, $g(x, y) = 0$, then we can say that the critical points satisfy the *Lagrange condition* $\nabla f_P = \lambda \nabla g_P$. This is equivalent to the *Lagrange equations*:

$$f_x(x, y) = \lambda g_x(x, y)$$

$$f_y(x, y) = \lambda g_y(x, y)$$

For functions in 3 variables, use 2 constraints: $g(x, y, z) = 0$ and $h(x, y, z) = 0$, then use two Lagrange multipliers:

$$\nabla f = \lambda \nabla g + \mu \nabla h$$

Multiple integration

Multiple integration can be computed in any order, for example:

$$\iint f(x, y) dx dy = \iint f(x, y) dy dx$$

Just integrate them as if they were single integrals, multiple times. This property is known as *Fubini's theorem*.

Riemann Sums in Multiple Variables

A *Riemann sum* for $f(x, y)$ on $\mathcal{R} = [a, b] \times [c, d]$ is a sum in the form:

$$S_{N,M} = \sum_{i=1}^N \sum_{j=1}^M f(P_{ij}) \Delta x_i \Delta y_j$$

Double integrals over complex regions

Assume that \mathcal{D} is a closed, bounded domain whose boundary is a simple closed curve that is either smooth or has a finite number of corners. The double integral is defined by:

$$\iint_{\mathcal{D}} f(x, y) dA = \iint_{\mathcal{D}} \tilde{f}(x, y) dA$$

where $\tilde{f}(x, y) = f(x, y)$ if $(x, y) \in \mathcal{D}$, otherwise $\tilde{f}(x, y) = 0$.

If \mathcal{D} is vertically simple, meaning $a \leq x \leq b$ and $g_1(x) \leq y \leq g_2(x)$, then evaluate:

$$\int_a^b \int_{g_1(x)}^{g_2(x)} f(x, y) dy dx$$

If \mathcal{D} is horizontally simple, meaning $h_1(y) \leq x \leq h_2(y)$ and $a < y < b$, then evaluate:

$$\int_a^b \int_{h_1(y)}^{h_2(y)} f(x, y) dx dy$$

Properties of Multiple Integration

If m is the minimum value and M is the maximum value of f on \mathcal{D} , then

$$m \cdot \text{area}(\mathcal{D}) \leq \iint_{\mathcal{D}} f(x, y) dA \leq M \cdot \text{area}(\mathcal{D})$$

If $z_1(x, y) \leq z_2(x, y) \forall (x, y) \in \mathcal{D}$, then the volume, V , of the solid region between the surfaces given by $z = z_1(x, y)$ and $z = z_2(x, y)$ over \mathcal{D} is

$$V = \iint_{\mathcal{D}} (z_2(x, y) - z_1(x, y)) dA$$

Mean value theorem

The mean value of f over \mathcal{D} , \bar{f} is

$$\bar{f} = \frac{1}{\text{area}(\mathcal{D})} \iint_{\mathcal{D}} f(x, y) dA = \frac{\iint_{\mathcal{D}} f(x, y) dA}{\iint_{\mathcal{D}} 1 dA}$$

If $f(x, y)$ is continuous and \mathcal{D} is closed, bounded, and connected, there exists a point P such that $f(P) = \bar{f}$.

Integration in other coordinate systems

Integration in polar coordinates:

$$\iint_{\mathcal{D}} f(x, y) dA = \int_{\theta_1}^{\theta_2} \int_{r_1}^{r_2} f(r \cos(\theta), r \sin(\theta)) r dr d\theta$$

Integration in cylindrical coordinates:

$$\iiint_{\mathcal{D}} f(x, y, z) dA = \int_{\theta_1}^{\theta_2} \int_{r_1}^{r_2} \int_{z_1(r, \theta)}^{z_2(r, \theta)} f(r \cos(\theta), r \sin(\theta), z) r dz dr d\theta$$

Integration in spherical coordinates

$$\iiint_{\mathcal{D}} f(x, y, z) dA = \int_{\theta_1}^{\theta_2} \int_{\phi_1}^{\phi_2} \int_{\rho_1}^{\rho_2} f(\rho \sin(\phi) \cos(\theta), \rho \sin(\phi) \sin(\theta), \rho \cos(\phi)) \rho^2 d\rho d\phi d\theta$$

Masses

For a mass with constant density, the center of mass coincides with the *centroids*, or geometric centers \bar{x} , \bar{y} , and \bar{z} , where

$$\begin{aligned} \langle \bar{x}, \bar{y}, \bar{z} \rangle &= \frac{1}{A} \iiint_{\mathcal{D}} \langle x, y, z \rangle dA \\ A &= \iiint_{\mathcal{D}} 1 dA \end{aligned}$$

For a mass with density function $\delta(x, y, z)$

$$\begin{aligned} \text{mass } M &= \iiint_{\mathcal{W}} \delta(x, y, z) dV \\ \text{moment } M_{yz} &= \iiint_{\mathcal{W}} x \delta(x, y, z) dV \\ \text{moment } M_{xz} &= \iiint_{\mathcal{W}} y \delta(x, y, z) dV \\ \text{moment } M_{xy} &= \iiint_{\mathcal{W}} z \delta(x, y, z) dV \\ \text{center of mass} &= \left\langle \frac{M_{yz}}{M}, \frac{M_{xz}}{M}, \frac{M_{xy}}{M} \right\rangle \end{aligned}$$

Probability

Random variables X and Y have the joint probability density function $p(x, y)$ if

$$\mathbb{P}(a \leq X \leq b; c \leq Y \leq d) = \int_{x=a}^b \int_{y=c}^d p(x, y) dx dy$$

A joint probability density function must satisfy $p(x, y) \geq 0$ and

$$\int_{x=-\infty}^{\infty} \int_{y=-\infty}^{\infty} p(x, y) dx dy = 1$$

Vector Fields

A *vector field* assigns a vector to every point in a domain. It is generally represented by a triple or double for vectors. For example, $\vec{F} = \langle F_1, F_2, F_3 \rangle$. We assume that each component is a smooth function on its domain.

The *del* or *nabla* operator, ∇ , defines 3 operations for vectors, *gradient*, *divergence*, and *curl*.

$$\begin{aligned} \text{grad}(f) &= \nabla f = \left\langle \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right\rangle \\ \text{div}(\vec{F}) &= \nabla \cdot \vec{F} = \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z} \\ \text{curl}(\vec{F}) &= \nabla \times \vec{F} = \left\langle \frac{\partial F_3}{\partial y} - \frac{\partial F_2}{\partial z}, \frac{\partial F_1}{\partial z} - \frac{\partial F_3}{\partial x}, \frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right\rangle \end{aligned}$$

Line Integrals

An *oriented curve* \mathcal{L} is a curve with a direction. The *arc-length differential* is $ds = ||\vec{r}'(t)|| dt$. The *scalar line integral* over \mathcal{L} with parameterization $\vec{r}(t)$ is:

$$\int_{\mathcal{L}} f(x, y, z) ds = \int_a^b f(\vec{r}(t)) ||\vec{r}'(t)|| dt$$

The *vector line integral* over \mathcal{L} with parameterization $\vec{r}(t)$ is:

$$\int_{\mathcal{L}} \vec{F} \cdot d\vec{r} = \int_a^b \vec{F}(\vec{r}(t)) \cdot \vec{r}'(t) dt$$

This is equivalent to

$$\int_{\mathcal{L}} \vec{F} \cdot d\vec{r} = \int_a^b F_1 dx + F_2 dy + F_3 dz$$

The parameterization of \mathcal{L} must be regular ($\vec{r}'(t) \neq \vec{0}$) and must trace \mathcal{L} in the positive direction.

Applications of Line Integrals

If $\rho(x, y, z)$ is the mass or charge density along \mathcal{L} , the total mass or charge is equal to the scalar line integral:

$$\int_{\mathcal{L}} \rho(x, y, z) ds$$

If \vec{F} is the force exerted on an object along a path \mathcal{L} , the work, W exerted on that object is equal to:

$$W = \int_{\mathcal{L}} \vec{F} \cdot d\vec{r}$$

The work performed *against* F is equal to:

$$- \int_{\mathcal{L}} \vec{F} \cdot d\vec{r}$$

Conservative Vector Fields

A vector field is considered *conservative* if $\vec{F} = \nabla f$. f is called a *potential function* for \vec{F} . Any two potential functions, f , differ only by a constant term (assuming an open, connected domain). A conservative vector field also satisfies the condition $\nabla \times \vec{F} = 0$.

A vector field \vec{F} on a domain \mathcal{D} is *path independent* if for any two points $P, Q \in \mathcal{D}$, where \mathcal{L}_1 and \mathcal{L}_2 are curves in \mathcal{D} from P to Q . A vector field \vec{F} on a domain \mathcal{D} is *path independent* if for any two points $P, Q \in \mathcal{D}$, where \mathcal{L}_1 and \mathcal{L}_2 are curves in \mathcal{D} from P to Q .

$$\int_{\mathcal{L}_1} \vec{F} \cdot d\vec{r} = \int_{\mathcal{L}_2} \vec{F} \cdot d\vec{r}$$

The *Fundamental Theorem for Conservative Vector Fields* states that, if $\vec{F} = \nabla f$, then, for any path from P to Q in the domain of \vec{F} :

$$\int_{\mathcal{L}} \vec{F} \cdot d\vec{r} = f(Q) - f(P)$$

Therefore, all conservative vector fields are path independent. The converse is also true: on an open, connected domain, a path independent vector field is conservative.

One special case of this is where \vec{r} is a closed path ($P = Q$):

$$\oint_{\mathcal{L}} \vec{F} \cdot d\vec{r} = 0$$

Parameterized Surfaces

A *parameterized surface* is a surface, \mathcal{S} , whose points are described in the form, where the *parameters* u and v vary in a domain \mathcal{D} in the uv -plane.

$$G(u, v) = (x(u, v), y(u, v), z(u, v))$$

The tangent (\vec{T}_u, \vec{T}_v) and normal (\vec{N}) vectors are defined as:

$$\vec{T}_u = \frac{\delta G}{\delta u} \quad \vec{T}_v = \frac{\delta G}{\delta v}$$

$$\vec{N} = \vec{N}(u, v) = \vec{T}_u \times \vec{T}_v$$

The parameterization is considered *regular* at (u, v) if $\vec{N}(u, v) \neq 0$.

Standard Parameterizations

The cylinder of radius R with the z -axis as the central axis:

$$\begin{aligned} G(\theta, z) &= (R \cos(\theta), R \sin(\theta), z) \\ \vec{N} &= \vec{T}_\theta \times \vec{T}_z = R \langle \cos(\theta), \sin(\theta), 0 \rangle \\ dS &= \|\vec{N}\| d\theta dz = R d\theta dz \end{aligned}$$

The sphere of radius R , centered at the origin:

$$\begin{aligned} G(\theta, \phi) &= (R \cos(\theta) \sin(\phi), R \sin(\theta) \sin(\phi), R \cos(\phi)) \\ \vec{e}_r &= \langle \cos(\theta) \sin(\phi), \sin(\theta) \sin(\phi), \cos(\phi) \rangle \\ \vec{N} &= \vec{T}_\phi \times \vec{T}_\theta = (R^2 \sin(\phi)) \vec{e}_r \\ dS &= \|\vec{N}\| dx dy = R^2 \sin(\phi) d\phi d\theta \end{aligned}$$

Graph of $z = g(x, y)$

$$\begin{aligned} G(x, y) &= (x, y, g(x, y)) \\ \vec{N} &= \vec{T}_x \times \vec{T}_y = \langle -g_x, -g_y, 1 \rangle \\ dS &= \|\vec{N}\| dx dy = \sqrt{1 + g_x^2 + g_y^2} dx dy \end{aligned}$$

Surface Integrals

Surface integrals are, well, integrals over surfaces. To compute a surface integral, we first take the parameterized surface G , and compute:

$$\iint_{\mathcal{S}} f(x, y, z) dS = \iint_{\mathcal{D}} f(G(u, v)) \|\vec{N}(u, v)\| du dv$$

Area computations

The area of a surface can be computed using a surface integral:

$$\text{area}(\mathcal{S}) = \iint_{\mathcal{D}} \|\vec{N}(u, v)\| du dv$$

The quantity $\|\vec{N}\|$ is the *area scaling factor*. For a small region \mathcal{D} on the uv -plane, and $\mathcal{S} = G(\mathcal{D})$, for any sample point (u_0, v_0) on \mathcal{D} ,

$$\text{area}(\mathcal{S}) \approx \|\vec{N}(u_0, v_0)\| \text{area}(\mathcal{D})$$

Surface integrals of Vector Fields

A surface \mathcal{S} is *oriented* if a varying unit vector $\vec{n}(P)$ is specified for each point on \mathcal{S} .

The integral of a vector field F over the oriented surface \mathcal{S} is defined as the surface integral of the normal component $\vec{F} \cdot \vec{n}$ over \mathcal{S} . Vector surface integrals are computed using the formula

$$\begin{aligned} \iint_{\mathcal{S}} \vec{F} \cdot \vec{n} dS &= \iint_{\mathcal{S}} \vec{F} \cdot d\vec{S} \\ &= \iint_{\mathcal{D}} \vec{F}(G(u, v)) \cdot \vec{N}(u, v) du dv \end{aligned}$$

The surface integral over a vector field is called the *flux* of \vec{F} through \mathcal{S} . If \vec{F} is the flow rate of a fluid, the flux of \vec{F} through \mathcal{S} is the amount of fluid that flows through \mathcal{S} per unit time.

Green's Theorem

There are two notations for the line integral of a vector field on a plane:

$$\int_{\mathcal{L}} \vec{F} \cdot d\vec{r} = \int_{\mathcal{L}} F_1 dx + F_2 dy$$

$\delta\mathcal{D}$ represents the oriented boundary of \mathcal{D} . Green's theorem states:

$$\oint_{\delta\mathcal{D}} F_1 dx + F_2 dy = \iint_{\mathcal{D}} \left(\frac{\delta F_2}{\delta x} - \frac{\delta F_1}{\delta y} \right) dA$$

The area of \mathcal{D} enclosed by \mathcal{C} is therefore:

$$\text{area}(\mathcal{D}) = \oint_{\mathcal{C}} x dy = \oint_{\mathcal{C}} -y dx = \frac{1}{2} \oint_{\mathcal{C}} x dy - y dx$$

Circulation form of Green's theorem:

$$\oint_{\delta\mathcal{D}} \vec{F} \cdot d\vec{r} = \iint_{\mathcal{D}} \text{curl}_z(\vec{F}) dA$$

Where $\text{curl}_z(\vec{F}) = \frac{\delta F_2}{\delta x} - \frac{\delta F_1}{\delta y}$. For a two-dimensional vector-field, the quantity $\text{curl}_z(\vec{F})$ can be interpreted as *circulation per unit area*. If \mathcal{C} is a small circle centered at P enclosing the domain \mathcal{D} ,

$$\text{curl}_z(\vec{F})(P) \approx \frac{1}{\text{area}(\mathcal{D})} \oint_{\mathcal{C}} \vec{F} \cdot d\vec{r}$$

Flux form of Green's theorem:

$$\oint_{\delta\mathcal{D}} \vec{F} \cdot \vec{n} ds = \iint_{\mathcal{D}} \text{div}(\vec{F}) dA$$

For a two dimensional vector field \vec{F} , $\text{div}(\vec{F})$ is interpreted as *outward flux per unit area*. If \mathcal{C} is a small circle centered at P with the enclosing domain \mathcal{D} , then

$$\text{div}(\vec{F})(P) \approx \frac{1}{\text{area}(\mathcal{D})} \oint_{\mathcal{C}} \vec{F} \cdot \vec{n} ds$$

Stoke's Theorem

Stoke's theorem is an extension of Green's theorem to 3 dimensions. Let \mathcal{S} be an oriented surface with the parameterization $G : \mathcal{D} \rightarrow \mathcal{S}$, where \mathcal{D} is a domain

in the plane bounded by smooth, simple closed curves, and G is 1-to-1 and regular.

$$\oint_{\delta\mathcal{S}} \vec{F} \cdot d\vec{r} = \iint_{\mathcal{S}} \text{curl}(\vec{F}) \cdot d\vec{S}$$

If \mathcal{S} is a closed surface, then

$$\iint_{\mathcal{S}} \text{curl}(\vec{F}) \cdot d\vec{S} = 0$$

Curl is interpreted as a vector that encodes circulation per unit area. Given any point P and a unit normal vector \vec{n} , where \mathcal{C} is a small circle centered at P in the plane through P with normal vector \vec{n} , \mathcal{D} is the region enclosed by \mathcal{C} , and θ is the angle between $\text{curl}(\vec{F})(P)$ and \vec{n} .

$$\begin{aligned} \text{curl}(\vec{F})(P) \cdot \vec{n} &= \|\text{curl}(\vec{F})(P)\| \cos(\theta) \\ &\approx \frac{1}{\text{area}(\mathcal{D})} \oint_{\mathcal{C}} \vec{F} \cdot d\vec{r} \end{aligned}$$

Divergence Theorem

If \mathcal{W} is a region in \mathbb{R}^3 whose boundary $\delta\mathcal{W}$ is a surface oriented by the normal vectors pointing outside \mathcal{W} , then:

$$\iint_{\delta\mathcal{W}} \vec{F} \cdot d\vec{S} = \iiint_{\mathcal{W}} \text{div}(\vec{F}) dV$$

It follows that if $\text{div}(\vec{F}) = 0$, then \vec{F} has zero flux through the boundary $\delta\mathcal{W}$ for any \mathcal{W} on the domain of \vec{F} . Divergence can be interpreted as flux per unit volume, which means that the flux through a small closed surface containing a point P is approximately equal to $\text{div}(\vec{F})(P)$ times the enclosed volume.

Vector Field Operations

$$\begin{array}{ccccccc} f & \xrightarrow{\nabla} & \vec{F} & \xrightarrow{\text{curl}} & \vec{G} & \xrightarrow{\text{div}} & g \\ \text{Function} & & \text{Vector field} & & \text{Vector field} & & \text{Function} \end{array}$$

$$\text{curl}(\nabla f) = 0 \quad \text{div}(\text{curl}(\vec{F})) = 0$$

The inverse-square field $\vec{F}_{\text{IS}} = e_r/r^2$, defined $\forall r \neq 0$ satisfies $\text{div}(\vec{F}_{\text{IS}}) = 0$. The flux of \vec{F}_{IS} through a closed surface \mathcal{S} is 4π if \mathcal{S} contains the origin, and 0 otherwise.